

Recently formed age-disparate partnerships are associated with elevated HIV-incidence among young women in South Africa

Brendan Maughan-Brown^a, Atheendar Venkataramani^b,
 Ayesha B.M. Kharsany^c, Sean Beckett^d, Kaymarlin Govender^d,
 Lara Lewis^c, Cherie Cawood^e, David Khanyile^e and Gavin George^d

Objective: Cross-sectional and cohort studies draw different conclusions on whether age-disparate partnerships increase HIV-acquisition risk for young women. We investigated whether age-disparities were associated with HIV-infection risk early in relationships. This could result in the exclusion of women who seroconverted during high-risk age-disparate partnerships from cohort studies of HIV incidence – which exclude HIV-positive women – and explain null findings in these studies.

Design: Prospective cohort study.

Methods: We used data on 15–24-year-old, HIV-negative women in heterosexual partnerships ($N = 830$) in KwaZulu-Natal, South Africa. The association between age-disparate partnering (i.e., male partner ≥ 5 years older) and subsequent HIV seroconversion was assessed using Cox hazard models. We examined heterogeneity in HIV-acquisition risk by duration of partnership (defined by quartiles) at cohort enrolment.

Results: During 1139 person-years (mean: 1.4 years) of follow-up, 54 (6.5%) women seroconverted, a weighted HIV-incidence estimate of 4.41/100 person-years [95% confidence interval (CI): 3.30–6.06]. HIV-acquisition risk did not differ significantly between women in age-disparate vs. age-similar partnerships (adjusted hazard ratios: 1.10, 95% CI: 0.55–2.21). However, for women in the shortest partnership quartile (< 1.09 years) at baseline, risk of HIV seroconversion was higher for women in age-disparate partnerships (adjusted hazard ratios: 3.13, 95% CI: 1.02–9.65, $P = 0.047$). HIV acquisition was not statistically different by partnership type among women in longer partnerships.

Conclusion: The association between age-disparate partnerships and HIV-acquisition risk is evident early in young women's relationships. Results provide a potential explanation for null findings in cohort studies, whose research designs may exclude women in such partnerships, and affirms the elevated risk of HIV acquisition for young women in age-disparate relationships. Copyright © 2019 Wolters Kluwer Health, Inc. All rights reserved.

AIDS 2020, **34**:149–154

^aSouthern Africa Labour and Development Research Unit (SALDRU), School of Economics, University of Cape Town, Middle Campus, Cape Town, South Africa, ^bDepartment of Medical Ethics and Health Policy, Perelman School of Medicine, University of Pennsylvania, Philadelphia, USA, ^cCentre for the AIDS Programme of Research in South Africa (CAPRISA), ^dHealth Economics and HIV and AIDS Research Division (HEARD), University of KwaZulu-Natal, Durban, and ^eEpicentre AIDS Risk Management (Pty) Limited, Sandton, South Africa.

Correspondence to Brendan Maughan-Brown, PhD, Southern Africa Labour and Development Research Unit (SALDRU), School of Economics Building, University of Cape Town, Middle Campus, Private Bag, Rondebosch, 7701 Cape Town, South Africa. Tel: +27 21 650 5695; fax: +27 21 650 5697; e-mail: brendan.maughanbrown@gmail.com

Received: 27 March 2019; revised: 18 July 2019; accepted: 19 July 2019.

DOI:10.1097/QAD.0000000000002362

Keywords: age-disparate sexual relationships, HIV acquisition, HIV incidence, HIV-infection risk, intergenerational partners, partner age difference, South Africa

Introduction

HIV-infection risk remains high among young women in sub-Saharan Africa. In 2016, 26% of new HIV infections in Eastern and Southern Africa occurred among 15–24-year-old women, who make up only 10% of the population [1]. In South Africa alone, approximately 2000 young women were infected with HIV every week in 2016 [2].

Evidence is currently mixed on age-disparate partnerships – those in which the male partner is 5 or more years older – as an HIV-infection risk factor for young women. Evidence from several sources indicates that partnering with older men increases risk of HIV infection for young women [3–9]. This evidence aligns with epidemiological data: older male partners are more likely to be HIV infected with an unsuppressed viral load than younger partners [6]; and behavioural data: condomless sex [10–13], transactional sex [13], higher frequency of sex [14–16], and concurrent sexual partnering [17], are more prevalent in age-disparate partnerships.

In contrast, three longitudinal studies in South Africa have shown no association between age-disparate partnerships and HIV-acquisition among young women [18–20]. These cohort studies have brought into question the efficacy of HIV-prevention interventions targeting age-disparate partnerships [21]. However, a potential source of bias in longitudinal studies is selection into prospective cohorts. Cohort studies of HIV incidence exclude HIV-positive individuals at baseline. As age-disparate partnerships are characterized by significantly greater sexual risk behaviour, one would expect HIV transmission to occur more rapidly within serodiscordant age-disparate partnerships than in age-similar partnerships. Eligibility criteria for cohort studies would consequently result in the exclusion of a greater proportion of women already infected within age-disparate partnerships, and thus in relatively fewer HIV-negative women remaining at risk of HIV infection within these partnerships. That is, the effect of age-disparities in increasing HIV acquisition for young women may have occurred prior to cohort selection in many cases, as might seem to be indicated by positive associations in cross-sectional studies.

The potential bias from selection effects of this nature in cohort studies would vary on the basis of the length of women's relationships at study eligibility assessment, with the smallest effects among women in newly formed relationships. We used prospective cohort data on 15–24-year-old HIV-negative women to assess heterogeneity in the relationship between age-disparate partnering and HIV seroconversion by partnership duration at cohort

enrolment. The hypothesis was that age-disparate partnerships would be associated with an increased risk of HIV acquisition within more recent partnerships, with weaker associations for partnerships of longer duration.

Methods

We used longitudinal data on participants testing HIV-negative during the 2015/2016 survey of the HIV Incidence Provincial Surveillance System, conducted from July 2015 to June 2016 in two subdistricts [Vulindlela (rural) and Greater Edendale (periurban)] in KwaZulu-Natal, South Africa [22]. This province has the highest HIV prevalence in South Africa [23], with HIV prevalence of approximately 44% among 15–49-year-old women in the study region [24]. Baseline survey participants ($N = 10\,236$; 15–49 years old) were selected using a two-stage random sampling of enumeration areas and then households, and then random selection of one individual per household. Cohort follow-up was conducted from November 2016 to August 2017. At baseline and follow-up, venous blood samples were collected from all participants for HIV-antibody testing, and face-to-face questionnaires were administered.

Laboratory HIV testing was done using fourth-generation HIV enzyme immunoassays to test for HIV antibodies and antigens using enzyme Biomerieux Vironostika Uniform II Antigen/Antibody Microelisa system (BioMérieux, Marcy-l'Étoile, France) and HIV 1/2 Combi Roche Elecys (Roche Diagnostics, Penzberg, Germany). Positive tests were confirmed with a Western blot (Biorad assay, Bio-Rad Laboratories, Redmond, Washington, USA).

The primary outcome was HIV seroconversion. The date of seroconversion was assumed to be the midpoint between the baseline and follow-up survey dates. The exposure of interest was the difference in age between the participant and her male partner at enrolment. Data were collected on participants' three most recent sexual partners (within the previous 5 years). Two measures of partner age differences were created using data on current (i.e., had not ended) heterosexual partnerships at baseline. First, a binary variable to identify age-disparate partnerships. Second, a continuous variable of the difference in age between partners (number of years the male partner was older), left censored at zero. For women in multiple, heterosexual partnerships at baseline ($n = 32$; 4%), data on the most recent sexual partner were used.

Cox proportional hazard models, as in previous cohort studies [18,19], were used to assess the relationship between HIV acquisition and each measure of partner age difference. Multivariable models included potential sociodemographic and behavioural confounders: participant age; education (secondary school completion); area of residence (urban vs. rural); concurrent sexual partnerships (i.e., multiple, heterosexual partnerships at baseline); relationship duration (quartiles); alcohol consumption; and household income. Relationship duration was categorized into quartiles to allow for nonlinear associations with HIV risk, which may arise as a result of reproduction decision-making changes during relationships.

To assess heterogenous effects of age-disparity on HIV acquisition by partnership duration, we included an interaction term: age-disparate partnerships (0/1) multiplied by partnership duration quartiles. Quartiles were used to create a category (quartile 1) of individuals in partnerships of less than 1 year, our a-priori definition of a recently established relationship. As this choice was arbitrary, we tested the robustness of results using partnership duration tertiles, and using quintiles of partnership duration. All models were estimated in Stata, version 15 (StataCorp, College Station, Texas, USA) with cluster-robust standard error (SE) estimation, using enumeration areas as the clustering unit, and applied survey weights.

Results

A total of 1861 women aged 15–24 years were eligible for this cohort study, and 1411 (76%) enrolled [refer to Fig. S1, <http://links.lww.com/QAD/B534>, Supplemental

Digital Content (SDC) 1]. The analytical sample ($N=830$) excluded 567 women not in a sexual partnership at baseline and 14 women in same sex partnerships. Participants were followed for a total of 1139 person-years of follow-up (mean: 1.4 years). Days to follow-up ranged from 314 to 777, with a mean of 518 (SD 87) and a median of 501.

Our sample (refer to Table S1, <http://links.lww.com/QAD/B534>, SDC 1) comprised Black African participants, predominately 20–24 years old (65%), with a mean age of 20.5 years. 27.3% of women [95% confidence interval (CI): 23.5–31.0] were in an age-disparate partnership at baseline. The mean partnership duration was 2.95 years [SE: 0.097; range: 0–15.59 years (outlier of 23.43 years excluded)], with 23% of partnerships starting in the year preceding the baseline.

Fifty-four (6.5%) women seroconverted between baseline and endline (4.41 per 100 person-years, 95% CI: 3.30–6.06) (Table 1). The difference in crude HIV incidence between women in age-disparate and age-similar partnerships was small and not statistically significant (4.85 vs. 4.25, $P=0.708$). The adjusted hazard ratios (aHR) for the association between HIV acquisition and age-disparate partnering (aHR: 1.10; 95% CI: 0.55–2.21) and partnership age differences in years (aHR: 1.04, 95% CI: 0.94–1.14) were small and not statistically significant.

Substantial differences in crude HIV incidence were evident between women in age-similar and age-disparate partnerships by partnership duration (refer to Table S3, <http://links.lww.com/QAD/B534>, SDC1). Among partnerships of less than 1.09 years duration (quartile 1), HIV incidence was approximately three times higher

Table 1. Association between HIV acquisition and age differences within partnerships at baseline among 15–24-year-old women in KwaZulu-Natal, South Africa (2015–2017).

	Unweighted data		Weighted data		
	HIV infections/ person-years	HIV incidence (per 100 person-years) (95% CI)	HIV incidence (per 100 person-years) (95% CI)	Unadjusted Cox proportional hazard model HR (95% CI)	Adjusted Cox proportional hazard model ^a aHR (95% CI)
All women (15–24 years old)	54/1138.69	4.74 (3.63–6.19)	4.41 (3.29–6.06)		
Independent variable 1					
Age-disparate partnership					
No	36/830.83	4.33 (3.13–6.01)	4.25 (2.97–6.29)	Ref	Ref
Yes	18/307.87	5.85 (3.69–9.28)	4.85 (2.91–8.67)	1.14 (0.58–2.25)	1.10 (0.55–2.21)
Independent variable 2					
Partnership age-difference (in years, 0–27)				1.04 (0.95–1.14)	1.04 (0.94–1.14)

Refer to Table S2, <http://links.lww.com/QAD/B534>, Supplemental Digital Content 1, for the full multivariable Cox proportional hazard models of HIV acquisition. aHR, adjusted hazard ratio; CI, confidence interval; HR, hazard ratio.

^aMultivariable Cox proportional hazard model controlling for participant age (years); education [completion of secondary schooling (grade 12) or not]; area of residence (urban vs. rural); relationship duration (quartiles); concurrent sexual partner at baseline (no/yes); household income ($\leq R2500$, $R2501–6000$, $>R6000$); and alcohol use (whether the respondent drinks alcohol or not).




Fig. 1. Hazard ratios for the association between HIV-acquisition and age-disparate partnerships (reported at baseline) among 15–24-year-old women ($N = 830$), by partnership duration (quartiles). Hazard ratios estimated using Cox proportional hazard models with interactions between age-disparate partnerships (0/1) and relationship duration quartiles. [^]Adjusted models included the following control variables: participant age (years); education [completion of secondary schooling (grade 12) or not]; area of residence (urban vs. rural); concurrent sexual partner at baseline (no/yes); household income (\leq R2500, R2501–6000, >R6000); and alcohol use (whether the respondent drinks alcohol/not).

for women in age-disparate partnerships compared with women with similar-aged partners (11.83 vs. 3.59 per 100 person-years). In contrast, HIV incidence was lower among women in age-disparate partnerships for each of the other partnership duration quartiles. Among age-disparate partnerships, 44% (8/18) of HIV infections occurred in those that began the relationship 1.08 years, or fewer, prior to the baseline survey.

Figure 1 presents hazard ratios for each of the partnership duration quartiles, calculated using Cox proportional hazard models with interaction terms for age-disparity and each partnership duration quartile (for model coefficients, refer to Table S4, <http://links.lww.com/QAD/B534>, SDC1). Among women reporting the shortest 25% of partnerships at baseline, those in age-disparate partnerships were significantly more likely to seroconvert (aHR: 3.13, 95% CI: 1.02–9.65, $P = 0.047$). Note, the mean duration of partnerships in quartile one was similar across partnership type (age-disparate: 0.55

years, age-similar: 0.61 years; $P = 0.197$). HIV acquisition was not statistically different by partnership type among women in longer-length partnerships.

Sensitivity analyses showed that heterogeneous age-disparate effects by partnership duration were not sensitive to adjustment of the partnership duration measure to tertiles, or to quintiles (refer to Figs. S2 and S3, <http://links.lww.com/QAD/B534>, SDC1). Additional analysis was conducted among a different sample in newly established relationships: 15–24-year-old women ($N = 209$) who were not in a sexual partnership at baseline but reported a sexual partner in the 12 months prior to the follow-up survey. This analysis (refer to Table S5, <http://links.lww.com/QAD/B534>, SDC1) also found a positive association between HIV acquisition and age-disparate partnerships (aHR: 2.71, 95% CI: 0.69–10.61, $P = 0.150$) and years age-difference between partners (aHR: 1.22, 95% CI: 1.02–1.46, $P = 0.031$).

Discussion

Recent cohort studies have found no association between age-disparate partnering and HIV-acquisition among women in South Africa [18–20]. These findings have led policymakers to question whether programmes to mitigate risks from age-disparate partnerships are an effective HIV-prevention strategy for young women. Our results challenge this interpretation. Specifically, we found that a null relationship between age-disparate partnering and HIV incidence masked substantial heterogeneity in HIV-infection rates by length of partnership. Within recently formed partnerships at cohort enrolment, HIV seroconversion was significantly more likely for women with age-disparate partners.

Study results suggest that the null findings in previous cohort studies of HIV incidence [18–20] may have resulted from analytical designs that excluded HIV-positive women. A large effect of age-disparity on HIV acquisition early in partnerships would result in a high proportion of *high-risk* age-disparate partnerships being excluded from cohort studies. This would lead to the selection of a sample comprised predominantly of individuals in lower risk partnerships, which would bias estimates of the impact of age-disparate partnerships on HIV incidence towards the null, and/or make it difficult to detect meaningful associations due to loss of statistical power. Selection effects in particular may provide a potential explanation for why, unlike cohort studies, cross-sectional studies typically do find a positive association between age-disparities and HIV positivity. Cross-sectional studies, by design, do not exclude HIV-positive women.

We acknowledge several study limitations. Both incomplete reporting of sexual partners, which is likely in surveys [25], and misclassification of whether a partnership had ended might have led to selection bias. Self-reported partner age [26], and partnership duration might also be measured imperfectly. Social desirability bias might have increased the likelihood of underreporting male partners' age in the region, given a recent campaign discouraging partnerships with older men. In addition, our study was conducted in a relatively small geographic region, which creates uncertainty about the external validity of the results. Finally, small sample bias might have influenced our results.

In conclusion, our study provides evidence of a significantly elevated HIV-acquisition risk among young women in more recently established age-disparate partnerships. This finding provides a plausible explanation for the lack of association between age-disparities and HIV incidence found in previous cohort studies in South Africa. Our results provide further evidence that targeting risks associated with age-disparate partnerships may be effective in reducing HIV-incidence rates among young women.

Acknowledgements

We thank all of the participants of the HIV Incidence Provincial Surveillance System (HIPSS), as well as HIPSS coinvestigators and members of the HIPSS study team from the following organizations: Epicentre, CAPRISA, HEARD, NICD, and CDC. We also thank the HIPSS collaborating partners: The National Department of Health, Provincial KwaZulu-Natal Department of Health, uMgungundlovu Health District, the uMgungundlovu District AIDS Council, local municipal and traditional leaders, and community members for all their support throughout the HIPSS study. We are extremely grateful to Kassahun Ayalew for valuable feedback on previous versions of this article.

The HIV Incidence Provincial Surveillance System (HIPSS) is funded by a cooperative agreement (3U2GGH000372) between Epicentre and the US Centers for Disease Control and Prevention (CDC). Support was provided to B.M.B. by the National Research Foundation, South Africa, through the Research Career Advancement Fellowship. A.B.M.K. is supported by a joint South Africa – US Program for Collaborative Biomedical Research, National Institutes of Health grant (R01HD083343). The content, findings, and conclusions in this article are those of the author(s) and do not necessarily represent the official views of the CDC, or any other funder.

Authors' contributions: A.B.M.K. is the principal investigator of the HIV Incidence Provincial Surveillance System (HIPSS). A.B.M.K., C.C., and D.K., were responsible for HIPSS study operations and quality assurance. B.M.B., A.V., A.B.M.K., and G.G. were responsible for the concept of the analysis for this article. All authors contributed to the interpretation of the data, and article preparation; all authors approved the final version of the report.

Conflicts of interest

There are no conflicts of interest.

References

1. UNAIDS. *Ending AIDS: progress towards the 90-90-90 targets*. Geneva: UNAIDS; 2017.
2. South African National AIDS Council. *Let our actions count*. Pretoria: Republic of South Africa National Department of Health; 2017. <http://nsp.sanac.org.za/wp-content/uploads/2017/01/Final-NSP-Documents.pdf>. [Accessed 18 January 2019].
3. Gregson S, Nyamukapa C, Garnett G, Mason P, Zhuwau T, Caraël M, *et al.* **Sexual mixing patterns and sex-differentials in teenage exposure to HIV infection in rural Zimbabwe**. *Lancet* 2002; **359**:1896–1903.
4. Pettifor AE, Kleinschmidt I, Levin J, Rees HV, MacPhail C, Madikizela-Hlongwa L, *et al.* **A community-based study to examine the effect of a youth HIV prevention intervention on young people aged 15–24 in South Africa: results of the baseline survey**. *Trop Med Int Health* 2005; **10**:971–980.
5. Evans M, Risher K, Zungu N, Shisana O, Moyo S, Celentano DD, *et al.* **Age-disparate sex and HIV risk for young women from 2002 to 2012 in South Africa**. *J Int AIDS Soc* 2016; **19**:21310.

6. Maughan-Brown B, George G, Beckett S, Evans M, Lewis L, Cawood C, *et al.* **HIV risk among adolescent girls and young women in age-disparate partnerships.** *J Acquir Immune Defic Syndr* 2018; **78**:155–162.
7. Kelly RJ, Gray RH, Sewankambo NK, Serwadda D, Wabwire-Mangen F, Lutalo T, *et al.* **Age differences in sexual partners and risk of HIV-1 infection in rural Uganda.** *J Acquir Immune Defic Syndr* 2003; **32**:446–451.
8. Schaefer R, Gregson S, Eaton JW, Mugurungi O, Rhead R, Takaruzza A, *et al.* **Age-disparate relationships and HIV incidence in adolescent girls and young women: evidence from Zimbabwe.** *AIDS* 2017; **31**:1461–1470.
9. de Oliveira T, Kharsany ABM, Gräf T, Cawood C, Khanyile D, Grobler A, *et al.* **Transmission networks and risk of HIV infection in KwaZulu-Natal, South Africa: a community-wide phylogenetic study.** *Lancet HIV* 2017; **4**:e41–e50.
10. Beauclair R, Kassinjee R, Temmerman M, Welte A, Delva W. **Age-disparate relationships and implications for STI transmission among young adults in Cape Town, South Africa.** *Eur J Contracept Reprod Health Care* 2012; **17**:30–39.
11. Volpe E, Hardie T, Cerulli C, Sommers M, Morrison-Beedy M. **What's age got to do with it? Partner age difference, power, intimate partner violence, and sexual risk in urban adolescents.** *J Interpers Violence* 2013; **28**:2068–2087.
12. Malema BW. **Determinants of condom use in Botswana: an empirical investigation of the role of gender.** *Botswana J Econ* 2012; **10**:59–78.
13. Maughan-Brown B, Evans M, George G. **Sexual behaviour of men and women within age-disparate partnerships in South Africa: implications for young women's HIV risk.** *PLoS One* 2016; **11**:e0159162.
14. Beauclair R, Dushoff J, Delva W. **Partner age differences and associated sexual risk behaviours among adolescent girls and young women in a cash transfer programme for schooling in Malawi.** *BMC Public Health* 2018; **18**:403.
15. Ritchwood TD, Hughes JP, Jennings L, MacPhail C, Williamson B, Selin A, *et al.* **Characteristics of age-discordant partnerships associated with HIV risk among young South African women (HPTN 068).** *J Acquir Immune Defic Syndr* 2016; **72**:423–429.
16. George G, Maughan-Brown B, Beckett S, Evans M, Cawood C, Khanyile D, *et al.* **Coital frequency and condom use in age-disparate partnerships involving women aged 15 to 24: evidence from a cross-sectional study in KwaZulu-Natal, South Africa.** *BMJ Open* 2019; **9**:e024362.
17. Maughan-Brown B, Kenyon C, Lurie MN. **Partner age differences and concurrency in South Africa: implications for HIV-infection risk among young women.** *AIDS Behav* 2014; **18**:2469–2476.
18. Harling G, Newell ML, Tanser F, Kawachi I, Subramanian SV, Barnighausen T. **Do age-disparate relationships drive HIV incidence in young women? Evidence from a population cohort in Rural KwaZulu-Natal, South Africa.** *J Acquir Immune Defic Syndr* 2014; **66**:433–451.
19. Balkus JE, Nair G, Montgomery ET, Mishra A, Palanee-Phillips T, Ramjee G, *et al.* **Age-disparate partnerships and risk of HIV-1 acquisition among South African women participating in the VOICE trial.** *J Acquir Immune Defic Syndr* 2015; **70**:212–217.
20. Street RA, Reddy T, Ramjee G. **The generational effect on age disparate partnerships and the risk for human immunodeficiency virus and sexually transmitted infections acquisition.** *Int J STD AIDS* 2016; **27**:746–752.
21. Geffen N. **Anti 'Sugar-daddy' campaign not supported by evidence.** *Health-e News*; 2014, <https://www.health-e.org.za/2014/03/10/antisugar-daddy-campaign-supported-evidence/>. [Accessed 29 May 2018].
22. Kharsany AB, Cawood C, Khanyile D, Grobler A, Mckinnon LR, Samsunder N, *et al.* **Strengthening HIV surveillance in the antiretroviral therapy era: rationale and design of a longitudinal study to monitor HIV prevalence and incidence in the uMgungundlovu District, KwaZulu-Natal, South Africa.** *BMC Public Health* 2015; **15**:1149.
23. Shisana O, Rehle T, Simbayi L, Zuma K, Jooste S, Zunga N, *et al.* **South African national HIV prevalence, incidence and behaviour survey 2012.** Cape Town: Human Sciences Research Council Press; 2014.
24. Grobler A, Cawood C, Khanyile D, Puren A, Kharsany AB. **Progress of UNAIDS 90–90–90 targets in a district in KwaZulu-Natal, South Africa, with high HIV burden, in the HIPSS study: a household-based complex multilevel community survey.** *Lancet HIV* 2017; **4**:e505–e513.
25. Maughan-Brown B, Venkataramani A. **Measuring concurrent partnerships: potential for underestimation in UNAIDS recommended method.** *AIDS* 2011; **25**:1549–1551.
26. Harling G, Tanser F, Mutevedzi T, Barnighausen T. **Assessing the validity of respondents' reports of their partners' ages in a rural South African population-based cohort.** *BMJ Open* 2015; **5**:e005638.